

Aristolochia yachangensis, a new species of Aristolochiaceae from limestone areas in Guangxi, China

Ya Jin Luo¹, Shi Dong Ni¹, Qiang Jiang¹, Bo Gao Huang¹, Yan Liu²,
Yu Song Huang²

¹ Management Center of Yachang Orchid National Nature Reserve, Baise, Guangxi, 533209, China ² Guangxi Key Laboratory of Functional Phytochemicals Research and Utilization, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and the Chinese Academy of Sciences, Guilin, Guangxi, 541006, China

Corresponding author: Yu Song Huang (huang-yusong@163.com); Yan Liu (gxibly@163.com)

Academic editor: Elton John de Lirio | Received 1 April 2020 | Accepted 11 June 2020 | Published 16 July 2020

Citation: Luo YJ, Ni SD, Jiang Q, Huang BG, Liu Y, Huang YS (2020) *Aristolochia yachangensis*, a new species of Aristolochiaceae from limestone areas in Guangxi, China. PhytoKeys 153: 49–61. <https://doi.org/10.3897/phytokeys.153.52796>

Abstract

Aristolochia yachangensis B.G.Huang, Yan Liu & Y.S.Huang, a new species from limestone areas in Guangxi, China, is described and illustrated. It is morphologically most similar to *A. fangchi* Y.C.Wu ex L.D.Chow & S.M.Hwang, *A. petelotii* O.C. Schmidt and *A. championii* Merr. & Chun in shape of leaf blade, anther, gynostemium and inflorescence on old woody stems. However, it can be easily distinguished from the latter by shape of inflorescence, length of upper and lower portions of perianth tube, colour of the limb and throat. A table and a key to distinguish the new species from other morphologically similar *Aristolochia* species are also provided.

Keywords

Aristolochia, limestone flora, new taxa, north-western Guangxi, taxonomy

Introduction

The genus *Aristolochia* L. (s. l.) contains 600 species and widely distributes in tropical, subtropical and temperate regions of the world (González 2012; Zhu et al. 2019c). Based on recent studies on molecular phylogeny, chromosome and morphology of *Aristolochia*, some researchers have suggested that an old genus *Isotrema* Raf. should be

reinstated to accommodate species of *Endodeca* Raf. and *Aristolochia* subgen. *Siphsia* (Duch.) O.C.Schmidt (Zhu et al. 2019a). However, many researchers still advise to use the name *Aristolochia* rather than *Isotrema* (Do et al. 2019; Peng et al. 2019; Cai et al. 2020). In this paper, we use the name *Aristolochia* to describe a new species, because the genus name *Isotrema* is still controversial.

Currently, there are more than 70 species of *Aristolochia* known from China, including many new species that have been described from Yunnan, Guangxi, Guangdong, Zhejiang and Hainan in recent years (Gong et al. 2018; Zhu et al. 2018, 2019b, 2019d; Li et al. 2019; Peng et al. 2019; Zhou et al. 2019). As one of the most biodiverse regions of China, Guangxi has 22 *Aristolochia* species (Peng et al. 2019; Zhu et al. 2019c), including *A. bambusifolia* C.F.Liang ex H.Q.Wen, *A. longlinensis* Yan Liu & L.Wu and *A. gongchengensis*, Y.S.Huang, Y.D.Peng & C.R.Lin, which are endemic in the region (Qin and Liu 2010; Huang et al. 2015; Wu et al. 2015).

During a fieldwork in Yachang Orchid National Nature Reserve of north-western Guangxi, China in April 2019, we discovered a special flowering plant of Aristolochiaceae and speculated that it might be a new species of *Aristolochia*, based on its flower structure. We investigated this species at the same location again and collected specimens of young capsules in May 2019. In order to obtain more detailed morphological data, we came back to the same location once again and collected specimens of mature capsules in July 2019. After consulting Flora of China (Hwang et al. 2003) and other relevant literature (Merrill and Chun 1940; Liang 1975; Chow and Huang 1975; Hwang 1981; Cheng et al. 1988; Ma 1989a, 1989b; Ma and Cheng 1989; Wen 1992; Liu and Deng 2009; Xu et al. 2011; Huang et al. 2013, 2015; Wu et al. 2013, 2015; Do et al. 2014, 2015a, 2015b, 2016, 2017, 2019; Huong et al. 2014; Zhu et al. 2015, 2016, 2017a, 2017b, 2018, 2019b, 2019d; Do and Nghiem 2017; Gong et al. 2018; Li et al. 2019; Peng et al. 2019; Zhou et al. 2019; Cai et al. 2020), as well as comparisons amongst this unknown species and its morphologically most similar species, we confirmed that this species was clearly different from the known *Aristolochia* species. Hence, it is here described and illustrated as a new species.

Material and methods

Field observations have been conducted in flowering and fruiting *phases* at the type locality more than once. Measurements and assessments of morphological characters of the new species were based on living plants in the wild and specimens gathered from the type locality. All specimens were deposited in the herbarium of Guangxi Institute of Botany (IBK), as well as the herbarium of Guangxi Botanical Garden of Medicinal Plants (GXMG). The comparisons amongst *Aristolochia yachengensis* B.G. Huang, Yan Liu & Y.S.Huang, *A. fangchi* Y.C.Wu ex L.D.Chow & S.M.Hwang, *A. petelotii* O.C.Schmidt and *A. championii* Merr. & Chun were based on the descriptions from herbarium specimens (including types) at CDBI, CSH, CZH, GXMG, GXMI, GZAC, GZTM, HEAC, HITBC, IBK, IBSC, K, KUN, NAS, PE, PEM, SM and protologues

(Schmidt 1933; Merrill and Chun 1940; Liang 1975). Images of type specimens and dried herbarium specimens were gathered from JSTOR Global Plants (<http://plants.jstor.org>), Chinese Virtual Herbarium Website (<http://www.cvh.ac.cn/>) and Sharing Platform of IBK (<http://www.gxib.cn/spIBK/>). The materials about current habitat status and threatened factors were recorded in field observations. The assessment of risk of extinction of the new species was based on the IUCN Red List of Threatened Species Categories and Criteria and Guidelines for Using the IUCN Red List Categories and Criteria (IUCN 2001; IUCN Standards and Petitions Committee 2019).

Taxonomic treatment

Aristolochia yachangensis B.G.Huang, Yan Liu & Y.S.Huang, sp. nov.

urn:lsid:ipni.org:names:77210596-1

Figures 1–3, 4A–D

Diagnosis. *Aristolochia yachangensis* is morphologically similar to *A. fangchi* Y.C.Wu ex L.D.Chow & S.M.Hwang, *A. petelotii* O.C.Schmidt and *A. championii* Merr. & Chun, but can be distinguished from them by stems irregularly striate, sparsely yellowish-brown pubescent or glabrous; leaf blade 1.5–3 cm wide; cymes on old woody stems; basal portion of perianth tube 2–3 cm long, shorter than the upper; limb yellow, with dark purple mural-like stripes; throat yellow; capsule ellipsoid. Detailed morphological comparisons amongst the four species of *A. yachangensis*, *A. championii*, *A. petelotii* and *A. fangchi* are summarised in Table 1.

Type. CHINA. Guangxi Zhuang Autonomous Region: Baise City, Leye County, Huaping Town, Zhongjing (Yachang Orchid National Nature Reserve), 24°49.367'N, 106°24.029'E, 1341 m a.s.l., 29 July 2019, Z. C. Lu et al. 20190729YC4141 (holotype: IBK!; isotypes: IBK!, GXMG!).

Description. Shrubs climbing. Stems terete, irregularly striate, sparsely yellowish-brown pubescent or glabrous. Branchlets densely yellowish-brown pubescent. Leaf blade leathery, lanceolate to elliptic-lanceolate or linear-lanceolate, 5–15 × 1.5–3 cm, apex narrowly acuminate, base rounded or broadly cuneate, margin entire, adaxially glabrous except the pubescent midnerve and lateral veins, abaxially shallowly yellowish-brown pubescent, basal veins 3, lateral veins 5–8 pairs, conspicuous on both surfaces; petiole 1–1.5 cm long, slightly distorted, densely yellowish-brown pubescent. Cymes on old woody stems, 1–5-flowered; pedicel 1–2 cm long, pendulous, densely yellowish-brown pubescent; bracteole ovate-lanceolate, ca. 4 × 2 mm, densely yellowish-brown pubescent; perianth tube horseshoe-shaped; basal portion of tube 2–2.5 × 0.6–1 cm, shorter than the upper part, near the base of inner dark purple, densely villous, outside of tube mauve, densely yellowish-brown pubescent; upper portion of tube 2.5–3 × 0.5–0.8 cm, inner yellow, with dark purple stripes; limb subrotund-peltate, 4–6 cm in diam., yellow, with dark purple mural-like stripes, abaxially densely brown pubescent, margin shallowly 3-lobed, lobes apex mucronate; throat suborbicu-

Table 1. Morphological comparisons of key characters amongst *Aristolochia yachengensis*, *A. fangchi*, *A. petelotii* and *A. championii*.

Characters	<i>A. yachengensis</i>	<i>A. fangchi</i>	<i>A. petelotii</i>	<i>A. championii</i>
Young stem	irregularly striate, sparsely yellowish-brown pubescent or glabrous	obscurely striate, brown villous	striate, densely spreading yellowish-brown villous	striate, densely yellowish-brown villous
Leaf blade	lanceolate to elliptic–lanceolate or linear–lanceolate, 5–15 × 1.5–3 cm, base rounded or widely cuneate, lateral veins 5–8 pairs	oblong to ovate–oblong, rarely ovate–lanceolate, 6–15 × 3–5.5 cm, base rounded or cordate, lateral veins 4–6 pairs	narrowly ovate, ovate–oblong or lanceolate–ovate, 14–22.5 × 7–13 cm, base shallowly cordate, lateral veins 4–6 pairs	lanceolate to elliptic–lanceolate or linear–lanceolate, 15–30 × 2–5 cm, base rounded or shallowly cordate, lateral veins 6–15 pairs
Pedicel	1–2 cm long, densely yellowish-brown pubescent	5–7 cm long, densely brown villous	4–4.5 cm long, densely brown villous	3–4 cm long, densely brown villous
Perianth tube	basal portion of tube 2–2.5 × 0.6–1 cm, shorter than the upper part, outside of tube mauve, densely yellowish-brown pubescent	basal portion of tube 4–5 × 1–1.5 cm, longer than the upper, outside of tube purple, with white blotches or not, densely villous	basal portion of tube 5–6.5 × 1–2 cm, longer than the upper, outside of tube pale-yellow or mauve, densely villous	basal portion of tube 5–7 × ca. 1.5 cm, longer than the upper, outside of tube mauve, densely villous
Limb	yellow, with dark purple mural–like stripes	dark purple, with white blotches	dark-purple, with white stripes	dark purple
Throat	yellow	white	milk-white mixed with black	yellow, with dark purple pots
Capsule	ellipsoid, 6–10 × 2.5–3.5 cm, glabrous	cylindrical, 5–10 × 3–5 cm, villous	narrowly ellipsoid, 10–15 × 5–8 cm, yellowish-brown villous	ellipsoid, 6–8 × ca. 3 cm, villous

lar, 0.5–1 cm in diam., yellow; anthers oblong, 2–4 × 1 mm, adnate to the gynostemium base, opposite to the lobes; ovary terete, ca. 1.5 × 0.3–0.5 cm, 6–angled, densely brown pubescent; gynostemium 3–lobed, margin glabrous and papillary. Capsule ellipsoid, 6–10 × 2.5–3.5 cm, 6–angled, glabrous.

Phenology. The new species was observed flowering from March to May and fruiting from June to August.

Etymology. The specific epithet is derived from the type locality, Yachang Orchid National Nature Reserve, Guangxi, China. The Chinese name is given as “雅长马兜铃”.

Distribution and habitat. At present, *Aristolochia yachangensis* was found only in Yachang Orchid National Nature Reserve of north-western Guangxi, China. It grows on limestone hillside at an elevation of ca. 1340 m. The slope direction is to the southwest, the slope is up to 40°, the tree layer is up to 15 m tall, the canopy cover is 70%, the shrub layer cover is 80% and the herb layer cover is 45%. Its associated species include *Quercus variabilis* Blume (Fagaceae), *Celtis sinensis* Pers. (Ulmaceae), *Platycarya longipes* Wu (Juglandaceae), *Toxicodendron succedaneum* (L.) Kuntze (Anacardiaceae), *Yua thomsonii* (Laws.) C.L.Li (Vitaceae), *Pteridium aquilinum* (L.) Kuhn var. *latiusculum* (Desv.) Underw. ex A.Heller (Pteridiaceae), *Miscanthus sinensis* Andersson (Gramineae) etc.




Figure 1. *Aristolochia yachangensis* B.G.Huang, Yan Liu & Y.S.Huang, sp. nov. **A** habit **B** flowering branch **C** flower (front view) **D** longitudinally dissected flower (showing the inside structure) **E** anthers and gynostemium (lateral view) **F** old phase of gynostemium (vertical view) **G** capsule. Drawn by Wenhong Lin (IBK).

Conservation status. Thus far, *Aristolochia yachangensis* has been found only from the type locality. The only subpopulation is located within a protected region and has seven individuals, including two mature ones. Based on the present study, its Extent of




Figure 2. *Aristolochia yachangensis* B.G.Huang, Yan Liu & Y.S.Huang, sp. nov. **A** habitat **B** flowering branch **C** flowers (front view) **D** flower (lateral view) **E** flower bud **F** longitudinally dissected flower (showing the inside structure) **G** longitudinally dissected flower (dorsal view) **H** old phase of gynostemium (vertical view) **I** old phase of anthers and gynostemium (lateral view) **J** ovary **K** young capsule **L** mature capsule. Photographed by Shuwan Li.




Figure 3. Holotype of *Aristolochia yachangensis* B.G.Huang, Yan Liu & Y.S.Huang, sp. nov. Z. C. Lu et al. 20190729YC4141(IBK).

Occurrence (EOO) is less than 100 km² and the known Area of Occupancy (AOO) is less than 0.5 km². According to Guidelines for Using the IUCN Red List Categories and Criteria (IUCN Standards and Petitions Committee 2019), *A. yachangensis* should




Figure 4. **A–D** *Aristolochia yachangensis* B.G.Huang, Yan Liu & Y.S.Huang, sp. nov. **A** habitat **B** inflorescence and flowers (lateral view) **C** flower (front view) **D** anthers and gynostemium **E–H** *A. championii* Merr. et Chun: **E** habitat **F** inflorescence and flower (front view) **G** flower (lateral view) **H** anthers and gynostemium **I–L** *A. fangchi* Y. C. Wu ex L. D. Chow et S. M. Hwang: **I** habitat **J** inflorescence and flower (lateral view) **K** flower (front view) **L** anthers and gynostemium **M–P** *A. petelotii* O. C. Schmidt: **M** habitat **N** inflorescence and flower (lateral view) **O** flower (front view) **P** anthers and gynostemium. Illustration by Wenhong Lin (based on the illustrations from Flora Reipublicae Popularis Sinicae).

be given a Vulnerable (VU) status, based on the criteria D2 of IUCN. As a newly-found species, however, it is probable that more subpopulations of *A. yachangensis* could be found in similar habitats of limestone areas of north-western Guangxi and southern Guizhou, China in the future.

Additional specimens examined (paratypes). CHINA. Guangxi Zhuang Autonomous Region: Baise City, Leye County, Huaping Town, Zhongjing (Yachang Orchid National Nature Reserve), 24°49.367'N, 106°24.029'E, 1341 m a.s.l., 21 April 2019, *Y. J. Luo & S. W. Li* 20190421001 (IBK); the same location, 17 May 2019, *Y. J. Luo et al.* YC4439 (IBK).

Discussion

Aristolochia yachangensis is unique in morphology. It is mostly similar to *A. fangchi*, *A. petelotii* and *A. championii*, but can be distinguished from all other *Aristolochia* species mainly based on the morphological characters of inflorescence, perianth tube, limb and throat. *A. yachangensis* can be further distinguished from morphologically-close species with the following key.

Key to *Aristolochia yachangensis* and morphologically-close species

- | | | |
|---|--|--------------------------|
| 1 | Limb adaxially papillate or upper papillate, lower smooth..... | 2 |
| – | Limb adaxially smooth..... | 3 |
| 2 | Basal portion of tube shorter than the upper; limb adaxially yellow, with dark purple stripes | 4 |
| – | Basal portion of tube longer than the upper; limb adaxially dark purple | |
| | <i>A. championii</i> | |
| 3 | Limb adaxially yellow | 5 |
| – | Limb adaxially dark purple or reddish-purple, sometimes with yellow or white blotches..... | 6 |
| 4 | Leaf blade narrowly ovate to ovate-oblong, base cordate; petiole 4–5 cm long; limb 3–4 cm in diam | <i>A. huanjiangensis</i> |
| – | Leaf blade lanceolate to elliptic-lanceolate or linear-lanceolate, base rounded or broadly cuneate; petiole 1–1.5 cm long; limb 4–6 cm in diam | |
| | <i>A. yachangensis</i> | |
| 5 | Leaf blade ovate to narrowly ovate; limb ca. 2.5 cm in diam.; lobes of gynostemium pubescent | <i>A. pilosistyla</i> |
| – | Leaf blade oblanceolate to lanceolate-elliptic; limb 4–6 cm in diam.; lobes of gynostemium glabrous..... | <i>A. versicolor</i> |
| 6 | Limb small, ca. 3 cm × 1.5–2 cm, adaxially without blotches.... | <i>A. fulvicoma</i> |
| – | Limb large, 4–13 cm in diam..... | 7 |

- 7 Leaf blade lanceolate–oblong or narrowly oblong, base narrowly auriculate, lateral veins 8–12; limb 8–13 cm in diam *A. westlandii*
- Leaf blade ovate, oblong or ovate-oblong, rarely ovate-lanceolate, base cordate or rounded; limb no more than 8 cm in diam **8**
- 8 Leaf blade base rounded, rarely cordate; limb dark purple, with pale yellowish blotches *A. fangchi*
- Leaf blade base cordate; limb dark purple or reddish-purple, with white blotches or pale yellowish, without blotches *A. petelotii*

Acknowledgements

The authors are very grateful to Mr. Wen-Hong Lin for preparing the illustration. We are also grateful to Mr. Shu-Wan Li, Mr. Ying Qin, Miss Hai-Ling Chen and Miss Zhao-Cen Lu for their assistance in fieldwork. This study was supported by the National Natural Science Foundation of China (grant no. 41661012), Special Funds for Local Science and Technology Development Guided by the Central Committee (grant no. ZY1949013), the Science & Technology Basic Resources Investigation Program of China (Grant No. 2017FY100100) and the Traditional Chinese medicine public health special project-The project of investigating and monitoring on the Chinese materia medica raw materials resources for national essential drugs ([2011]76) and Traditional Chinese Medicine industry research special project-Characteristic Chinese materia medica resources protection and utilization in representative regions of China (201207002).

References

- Cai L, He DM, Huang YS, Dao ZL (2020) *Aristolochia wenshanensis*, a new species of Aristolochiaceae from karst region in southeastern Yunnan, China. *Taiwania* 65: 41–46.
- Cheng CY, Yang CS, Hwang SM (1988) *Aristolochia* Linnaeus. In: Kiu HS, Ling YR (Eds) *Flora Reipublicae Popularis Sinicae*. Science Press, Beijing, 199–245.
- Chow LD, Huang SM (1975) *Aristolochia fangchi*, a new species of *Aristolochia* (Aristolochiaceae). *Zhiwu Fenlei Xuebao* 13: 108–110.
- Do TV, Nghiem TD (2017) Taxonomic notes on some *Aristolochia* species in Vietnam. *Taiwania* 62: 216–218.
- Do TV, Nghiem TD, Wanke S, Neinhuis C (2014) *Aristolochia quangbinhensis* (Aristolochiaceae), a new species from Central Vietnam. *PhytoKeys* 33: 51–59. <https://doi.org/10.3897/phytokeys.33.6094>
- Do TV, Luu TH, Wanke S, Neinhuis C (2015a) Three new species and three new records of *Aristolochia* subgenus *Siphisia* from Vietnam including a key to the Asian species. *Systematic Botany* 40(3): 671–691. <https://doi.org/10.1600/036364415X689140>

- Do TV, Neinhuis C, Wanke S (2015b) A new species of *Aristolochia* subgenus *Siphisia* (Aristolochiaceae) from central Vietnam. *Phytotaxa* 220(1): 69–76. <https://doi.org/10.11646/phytotaxa.220.1.6>
- Do TV, Wanke S, Neinhuis C (2016) *Aristolochia bidoupensis* sp. nov. from southern Vietnam. *Nordic Journal of Botany* 34(5): 513–516. <https://doi.org/10.1111/njb.01066>
- Do TV, Truong CQ, Huynh HTT (2017) *Aristolochia neinhuisii* (Aristolochiaceae), a new species from Vietnam. *Annales Botanici Fennici* 54(4–6): 203–208. <https://doi.org/10.5735/085.054.0602>
- Do TV, Vu TTH, Luu HT, Nguyen TT (2019) *Aristolochia nuichuaensis* (subg. *Siphisia*, Aristolochiaceae), a new species, an updated key and a checklist to the species of *Siphisia* in Vietnam. *Annales Botanici Fennici* 56(1–3): 107–113. <https://doi.org/10.5735/085.056.0116>
- Gong QB, Landrein S, Xi HC, Ma XD, Yang ZH, He KW, Shen JY (2018) *Aristolochia tongbi-guanensis*, a new species of Aristolochiaceae from Yunnan, China. *Taiwania* 63: 183–187.
- González F (2012) Florística y sistemática filogenética innecesariamente disyuntas: El caso de *Aristolochia*, *Euglypha* y *Holostylis* (Aristolochiaceae). *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 36: 193–202.
- Huang YS, Peng RC, Tan WN, Wei GF, Liu Y (2013) *Aristolochia mulunensis* (Aristolochiaceae), a new species from limestone areas in Guangxi, China. *Annales Botanici Fennici* 50(3): 175–178. <https://doi.org/10.5735/085.050.0308>
- Huang YS, Peng YD, Huang BY, Lv HZ, Lin CR (2015) *Aristolochia gongchengensis* (Aristolochiaceae), a new species from the limestone areas in Guangxi, China. *Annales Botanici Fennici* 52: 396–400. <https://doi.org/10.5735/085.052.0522>
- Huong NTT, Hai DV, Quang BH, Cuong NT, Kuang NS, Vu DQ, Ma JS (2014) *Aristolochia xuanlienensis*, a new species of Aristolochiaceae from Vietnam. *Phytotaxa* 188(3): 176–180. <https://doi.org/10.11646/phytotaxa.188.3.7>
- Hwang SM (1981) Materials for Chinese *Aristolochia*. *Zhiwu Fenlei Xuebao* 19: 222–231.
- Hwang SM, Kelly LM, Gilbert MG (2003) *Aristolochia* Linnaeus. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China*, Vol. 5. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 258–269.
- IUCN (2001) IUCN Red List Categories and Criteria, Version 3.1. Gland Switzerland and Cambridge, 18–20.
- Li RT, Wang ZW, Wang J, Zhu XX, Xu H (2019) *Isotrema sanyaense*, a new species of Aristolochiaceae from Hainan, China. *PhytoKeys* 128: 85–96. <https://doi.org/10.3897/phytokeys.128.35042>
- Liang CF (1975) The Aristolochiaceae of Kwangsi Flora. *Zhiwu Fenlei Xuebao* 13: 10–28.
- Liu ZW, Deng YF (2009) *Aristolochia wuana*, a new name in Chinese *Aristolochia* (Aristolochiaceae). *Novon* 19(3): 370–371. <https://doi.org/10.3417/2007151>
- Ma JS (1989a) A revision of *Aristolochia* Linn. from E. & S. Asia. *Zhiwu Fenlei Xuebao* 27: 321–364.
- Ma JS (1989b) A revision of *Aristolochia* from Yunnan. *Yunnan Zhi Wu Yan Jiu* 11: 321–323.
- Ma JS, Cheng JR (1989) New taxa of Chinese *Aristolochia* L. *Zhiwu Fenlei Xuebao* 27: 293–297.
- Merrill ED, Chun WY (1940) Additions to our knowledge of the Hainan flora. *Sunyatsenia* 5: 45–200.

- Peng YD, Gadagkar SR, Li J, Xie YY, Huang XY, Lu HZ, Huang BY, Yu LY (2019) *Aristolochia kechangensis* sp. nov. (Aristolochiaceae) from Guangxi, China. *Nordic Journal of Botany* 37(9): 1–7. <https://doi.org/10.1111/njb.02456>
- Qin HN, Liu Y (2010) A checklist of vascular plants of Guangxi. Science Press, Beijing, 82–83.
- Schmidt OC (1933) Beiträge zur Kenntnis der Aristolochiaceen IV. Repertorium Specierum Novarum Regni Vegetabilis 32(1–8): 95–96. <https://doi.org/10.1002/fedr.19330320104>
- Standards IUCN, Petitions Committee (2019) Guidelines for Using the IUCN Red List Categories and Criteria. Version 14. Prepared by the Standards and Petitions Committee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Wen HQ (1992) Species of *Aristolochia* of Guangxi. *Gaihaia* 12: 217–218.
- Wu L, Xu WB, Wei GF, Liu Y (2013) *Aristolochia huanjiangensis* (Aristolochiaceae), a new species from Guangxi, China. *Annales Botanici Fennici* 50(6): 413–416. <https://doi.org/10.5735/085.050.0608>
- Wu L, Xu WB, Huang YS, Liu Y (2015) *Aristolochia longlinensis* (Aristolochiaceae), a new species from Western Guangxi, China. *Novon* 23(4): 490–493. <https://doi.org/10.3417/2011105>
- Xu H, Li YD, Yang HJ, Chen HQ (2011) Two new species of *Aristolochia* (Aristolochiaceae) from Hainan Island, China. *Novon* 21(2): 285–289. <https://doi.org/10.3417/2009116>
- Zhou XX, Jiang GB, Zhu XX, Liu ZY, Huang Y, Wang GT, Wang RJ (2019) *Isotrema plagiostomum* (Aristolochiaceae), a new species from Guangdong, South China. *Phytotaxa* 405(4): 221–225. <https://doi.org/10.11646/phytotaxa.405.4.7>
- Zhu XX, Zhang L, Hua ZX, Chen GF, Liao S, Ma JS (2015) *Aristolochia weixiensis*, a new species of Aristolochiaceae from Yunnan, China. *Phytotaxa* 230(1): 54–60. <https://doi.org/10.11646/phytotaxa.230.1.4>
- Zhu XX, Liao S, Zhang L, Wang ZH, Du C, Ma JS (2016) The taxonomic revision of Asian *Aristolochia* (Aristolochiaceae) I: Confirmation and illustration of *A. austroszechuanica*, *A. faucimaculata* and *A. yunnanensis* var. *meionantha* from China. *Phytotaxa* 261(2): 137–146. <https://doi.org/10.11646/phytotaxa.261.2.3>
- Zhu XX, Liao S, Ma ZX, Xu B, Wang ZH, Wang Y, Ma JS (2017a) The taxonomic revision of Asian *Aristolochia* (Aristolochiaceae) III: Two new taxa of *Aristolochia* and morphological revision for the flower character of *A. obliqua* from Yunnan, China. *Phytotaxa* 332(3): 269–279. <https://doi.org/10.11646/phytotaxa.332.3.3>
- Zhu XX, Liao S, Sun ZP, Zhen AG, Ma JS (2017b) The taxonomic revision of Asian *Aristolochia* (Aristolochiaceae) II: Identities of *Aristolochia austroyunnanensis* and *A. dabieshanensis*, and *A. hyperxantha*-a new species from Zhejiang, China. *Phytotaxa* 313(1): 61–76. <https://doi.org/10.11646/phytotaxa.313.1.4>
- Zhu XX, Liao S, Liu JN, Zhang C, Ma JS (2018) The taxonomic revision of Asian *Aristolochia* (Aristolochiaceae) IV: Lectotypification of *A. caulialata*, with a new species from Yunnan, China – *A. pseudocaulialata*. *Phytotaxa* 364(1): 49–60. <https://doi.org/10.11646/phytotaxa.364.1.2>
- Zhu XX, Li XQ, Liao S, Du C, Wang Y, Wang ZH, Yan J, Zuo YJ, Ma JS (2019a) Reinstatement of *Isotrema*, a new generic delimitation of *Aristolochia* subgen. *Siphisia* (Aristolochiaceae). *Phytotaxa* 401(1): 1–23. <https://doi.org/10.11646/phytotaxa.401.1.1>

- Zhu XX, Li XQ, Liao S, Li GD, Ma JS (2019b) The taxonomic revision of Asian *Aristolochia* (Aristolochiaceae) V: Two new species from Yunnan, China. *PhytoKeys* 130: 93–106. <https://doi.org/10.3897/phytokeys.130.33933>
- Zhu XX, Wang J, Liao S, Ma JS (2019c) Synopsis of *Aristolochia* L. and *Isotrema* Raf. (Aristolochiaceae) in China. *Shengwu Duoyangxing* 27(10): 1143–1146. <https://doi.org/10.17520/biods.2019183>
- Zhu XX, Zheng HL, Wang J, Gao YQ, Ma JS (2019d) Taxonomic studies on the genus *Isotrema* (Aristolochiaceae) from China: 1. *I. cangshanense*, a new species from Yunnan. *PhytoKeys* 134: 115–124. <https://doi.org/10.3897/phytokeys.134.37243>